

Comment on

” Thermodynamics of a One- Dimensional Ideal Gas with fractional Exclusion Statistics”

Alain DASNIÈRES de VEIGY and Stéphane OUVRY¹

Division de Physique Théorique², IPN, Orsay Fr-91406

PACS numbers: 03.65.-w, 05.30.-d, 11.10.-z, 05.70.Ce

In [1], the conclusion was reached that, in the one-dimensional Calogero model, only the second virial coefficient is affected by the statistical parameter α , where α is related to the coupling constant κ/x_{ij}^2 of the Calogero interaction by $\kappa = \alpha(\alpha + 1)$ ($\alpha \rightarrow -g$ in their notation). We argue that it is not so, i.e. all virial coefficients are affected, if the thermodynamic limit is properly taken. In [1], the system is regularized at long distance by harmonically attracting the particles to the origin, with the harmonic pulsation $\omega \rightarrow 0$ in the thermodynamic limit. However, this limit should be taken with a particular prescription that will be detailed below. In fact, the correct cluster expansion was already conjectured by Sutherland in its original paper [2] on the thermodynamic of the Calogero model, by various means, in particular a harmonic regularization. On the other hand, the anyon model in the lowest Landau level (LLL) of an external B -field [3] (LLL-anyon model) shares some similarities with the Calogero model, if the anyonic statistical parameter $\alpha \in [-1, 0]$ modulo 2 (by convention, $eB > 0$ has been assumed). The link with Haldane’s statistics Hilbert space counting argument [4] is immediate, because of the diminution of available one-body quantum states in the LLL, due to the screening of the B field. In this regime, the groundstate is known, excited states have a gap, and the LLL-anyon model can be viewed as a one-dimensional model. In the sequel, we will treat simultaneously both models -for a review on the subject see [5].

In a harmonic well, the N -body spectrum of the Calogero or LLL-anyon models is

¹and LPTPE, Tour 12, Université Paris 6 / electronic e-mail: OUVRY@FRCPN11

²Unité de Recherche des Universités Paris 11 et Paris 6 associée au CNRS

$\sum_{\ell} n_{\ell} \epsilon_{\ell} - \frac{1}{2} N(N-1) \varpi \alpha$, where, for a given angular momentum l , the 1-body energy is $\epsilon_{\ell} = \epsilon_0 + \ell \varpi$. The LLL-anyon spectrum corresponds to $\varpi = \omega_t - \omega_c$ ($\omega_t = \sqrt{\omega_c^2 + \omega^2}$, $\omega_c = eB/(2m)$) and $\epsilon_0 = \omega_t$, whereas the Calogero spectrum corresponds to $\varpi = \omega$ and $\epsilon_0 = \frac{1}{2}\omega$. From the N -body partition function

$$Z_N^{\omega} = e^{\frac{1}{2}\beta N(N-1)\varpi\alpha} \prod_{n=1}^N \frac{e^{-\beta\epsilon_0}}{1 - e^{-n\beta\varpi}}, \quad (1)$$

one deduces the cluster expansion of the thermodynamical potential $\Omega^{\omega} \equiv -\ln \sum_N Z_N^{\omega} z^N = -\sum_n b_n^{\omega} z^n$. In the limit $\omega \rightarrow 0$

$$b_n^{\omega} = \frac{1}{\beta\varpi} \frac{e^{-n\beta\epsilon_0}}{n^2} \prod_{k=1}^{n-1} \frac{k + n\alpha}{k} \quad (2)$$

The thermodynamic limit prescription in (2) should be dictated by the requirement that when $\omega \rightarrow 0$, the thermodynamical potential of a system of d -dimensional harmonic oscillators coincide with those of a system of d -dimensional particles in a box of infinite volume $\Omega^V = -\sum_n b_n^V z^n$. Close to $\vec{r}$, one can approximate $\sum_i \frac{1}{2} m \omega^2 \vec{r}_i^2$ by $\frac{N}{2} m \omega^2 \vec{r}^2$. It follows that, for a system of particles confined in an infinitesimal volume $d^d \vec{r}$ around $\vec{r}$, the local thermodynamical potential in the presence of a small harmonic attraction can be approximated by replacing $z \rightarrow z e^{-\frac{1}{2}\beta m \omega^2 \vec{r}^2}$ and $V \rightarrow d^d \vec{r}$ in the infinite box thermodynamical potential. Since the thermodynamical potential is additive in the limit $\omega \rightarrow 0$, one has for the entire system (λ is the thermal wavelength)

$$\Omega^{\omega} = -\int \frac{d^d r}{V} \sum_{n=1}^{\infty} b_n^V \left(z e^{-\frac{1}{2}\beta m \omega^2 r^2} \right)^n = -\sum_{n=1}^{\infty} \frac{\lambda^d}{n^{d/2} (\beta\omega)^d V} b_n^V z^n \quad (3)$$

Thus one has $b_n^{\omega} \rightarrow b_n^V$, provided that the prescription

$$1/(n\beta^2\omega^2)^{d/2} \rightarrow V/\lambda^d \quad (4)$$

is enforced at each order z^n . One deduces the following results :

LLL-anyon model : The cluster coefficients are

$$b_n^V = \frac{V}{\lambda^2} \frac{e^{-n\beta\omega_c}}{n} \prod_{k=1}^{n-1} \frac{k + n\alpha}{k} \quad (5)$$

The thermodynamical potential reads $\Omega^V = -V\rho_L \ln y$, where y is solution of $y - ze^{-\beta\omega_c}y^{1+\alpha} = 1$. This is precisely because of the $d = 2$ prescription given in (4) that the Landau degeneracy $\rho_L V$ has factorized. The virial coefficients are $a_n = (-\frac{1}{\rho_L})^{n-1} \frac{1}{n} \{(1+\alpha)^n - \alpha^n\}$.

Calogero model : The cluster coefficients are

$$b_n^V = \frac{V}{\lambda^2} \frac{1}{n^{3/2}} \prod_{k=1}^{n-1} \frac{k+n\alpha}{k} \quad (6)$$

The thermodynamical potential is $\Omega^V = -\frac{V}{2\pi} \int dp \ln y$, where y is now solution of $y - ze^{-\beta p^2/2m}y^{1+\alpha} = 1$. The continuous one-dimensional energy spectrum $p^2/2m$ appears precisely because of the $d = 1$ prescription (4). The virial coefficients are all α dependent, $a_2 = -\sqrt{2}(\frac{1}{4} + \frac{\alpha}{2})$, $a_3 = \frac{1}{2} + 2\alpha + 2\alpha^2 - \sqrt{3}(\frac{2}{9} + \alpha + \alpha^2)$, et caetera. In [1], the inappropriate thermodynamic limit prescription $1/(\beta^2\omega^2)^{d/2} \rightarrow V/\lambda^d$ was taken, and a virial expansion interpolating between the $d = 2$ Bose and Fermi gases was obtained, where only the second virial coefficient is affected by α . It would be certainly satisfactory to find a $d = 2$ N -body Hamiltonian which would lead to such a simple virial expansion.

References

- [1] M. V. N. Murthy and R. Shankar, Phys. Rev. Lett. **73**, 3331 (1994).
- [2] B. Sutherland, J. Math. Phys. (N.Y) **12**, 251 (1971).
- [3] A. Dasnières de Veigy and S. Ouvry, Phys. Rev. Lett. **72**, 600 (1994).
- [4] F. D. M. Haldane, Phys. Rev. Lett. **67**, 937 (1991) ; Y.-S. Wu, Phys. Rev. Lett. **73**, 922 (1994); S. B. Isakov, Phys. Rev. Lett. **73**, 2150 (1994).
- [5] A. Dasnières de Veigy and S. Ouvry, Mod. Phys. Lett. B **9** (1995).